

ATH-1105, a small-molecule positive modulator of the neurotrophic HGF system, attenuates TDP-43 pathology in preclinical models of ALS



Sherif M. Reda, Andrée-Anne Berthiaume, Wei Wu, Kayla N. Kleist, Sharay E. Setti, Robert W. Taylor, Jewel L. Johnston, Kevin J. Church

Athira Pharma, Bothell, USA

Presented at ENCALs 2024; June 17-20; Stockholm, Sweden

CONCLUSIONS

- ATH-1105 promotes motor neuron survival and reduces cytoplasmic TDP-43 accumulation following glutamate injury**
- The neuroprotective effects of ATH-1105 are associated with improved autophagic function, which may promote the clearance of pathological TDP-43**
- ATH-1105 also promotes mitochondrial health and reduces GSK3 β activation, which may mitigate the generation of pathological TDP-43**
- In a TDP43-driven mouse model of ALS, ATH-1105 reduces TDP-43 pathology and improves neuromuscular function**


KEY TAKEAWAY

ATH-1105 addresses multiple modes of dysfunction associated with TDP-43 pathology, supporting its ongoing development as a potential treatment for ALS.

INTRODUCTION

- ALS pathology is associated with progressive motor neuron degeneration, glutamate excitotoxicity, mitochondrial dysfunction, oxidative stress, axonal degeneration, TDP-43 cytoplasmic accumulation, and motor neuron death^{1,2}
- TDP-43 is a highly conserved and ubiquitously expressed RNA/DNA-binding protein that has versatile functions in transcription, translation, and mRNA transport and stabilization³
- Nuclear depletion and cytoplasmic accumulation of TDP-43 is a critical component of ALS pathology that contributes to the progressive degeneration of motor neurons.
 - Up to ~97% of people with ALS exhibit TDP-43 proteinopathy⁴
- Cytoplasmic TDP-43 can manifest in various pathological forms (TDP-43 aggregates, TDP-43 fragments, and phosphorylated TDP-43), which are associated with several neurodegenerative mechanisms including autophagic impairment, mitochondrial dysfunction, and activation of GSK3 β .
- Promotion of neurotrophic HGF activity has been reported to have beneficial effects in preclinical models of ALS through its pleiotropic actions that can counteract various neurodegenerative mechanisms^{5,6}
- We sought to assess the impact of ATH-1105, a small molecule positive modulator of HGF, on TDP-43 pathology in preclinical models of ALS.

ATH-1105 is a positive modulator of the neurotrophic HGF system⁵



OBJECTIVE

To assess the impact of ATH-1105 on neurodegenerative mechanisms associated with TDP-43 pathology

METHODS

In vitro primary motor neuron glutamate toxicity assays

- Rat primary spinal motor neurons were harvested from E14 rat embryos and cultured for 13 days
- Cultures were pretreated for 15 minutes with vehicle (containing HGF 0.05 ng/ml) or ATH-1105, and then challenged with glutamate 5 μ M for 24 hours
- After 24 hours, immunofluorescence was used to assess the following metrics:
 - Neuronal survival (MAP2+ positive neurons)
 - Neurite network (total length of MAP2+ neurites in μ m)
 - Cytoplasmic TDP-43 (overlap area of TDP-43 and MAP2+ neurons in μ m²)
 - LC3+ autophagosomes (overlap area of LC3 and MAP2+ neurons in μ m²)
 - LAMP2+ lysosomes (overlap area of LAMP2 and MAP2+ neurons in μ m²)
 - Autophagosome-lysosome co-localization (overlap area of LC3-LAMP2 in μ m²)
 - Functional mitochondria (overlap of MitoTracker and MAP2+ neurons in μ m²)
 - Active caspase-3 (overlap between caspase-3 and MAP2+ neurons in μ m²)
- Western blot analysis was used to evaluate pTDP-43 (Ser409/410), pGSK3 β (Ser9), total GSK3 β , and GAPDH (reference protein)

In vivo assays in the Prp-TDP43^{A315T} mouse model of ALS

- 1-month-old male mice were used in the study. Mice were divided into four groups (n=10) and treated once daily for 2 months as follows:
 - WT mice treated with oral vehicle
 - Prp-TDP43^{A315T} mice treated with oral vehicle
 - Prp-TDP43^{A315T} mice treated with oral ATH-1105 10 mg/kg
 - Prp-TDP43^{A315T} mice treated with oral ATH-1105 20 mg/kg
- Quantification of pTDP-43 was performed in homogenized sciatic nerve using the AlphaLISA SureFire Ultra Human Phospho-TDP-43 (Ser409/410) detection kit
- Quantification of plasma NfL was performed in duplicate for each animal by NfL ELISA kit: Novus Biologica Ref. NBP2-80299
- Neuromuscular function was evaluated by measuring CMAP amplitude from the intrinsic foot muscles of anesthetized mice using steel-needle electrodes (MLA1302; AD Instruments)

RESULTS

Figure 1. ATH-1105 promotes motor neuron survival and reduces cytoplasmic TDP-43 following glutamate injury






Figure 2. ATH-1105 promotes phospho-inactivation of GSK3 β and reduces pTDP-43 following glutamate injury



Quantification of (A) Phospho-GSK3 β (Ser9) and (B) Phospho-TDP-43 (Ser409/410) 24h post glutamate challenge expressed as percentage of normal control (100%). Data presented as mean \pm SEM; n = 3-4. Statistical differences were determined by one-way ANOVA followed by Fisher's LSD test versus glutamate control. **p < 0.01, ***p < 0.001

Figure 3. ATH-1105 restores autophagic function following glutamate injury



Abbreviations: ALS, amyotrophic lateral sclerosis; AKT, protein kinase B; ANOVA, analysis of variance; CMAP, compound muscle action potential; ERK, extracellular signal regulated kinase; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; GSK3 β , glycogen synthase kinase-3 beta; HGF, hepatocyte growth factor; LAMP2, lysosomal membrane-associated protein 2; LC3, microtubule-associated protein 1A/1B light chain 3B; LSD, least significant difference; MAP2, microtubule-associated protein 2; NfL, neurofilament light; NMJ, neuromuscular junction; p, phosphorylation; pAKT, phosphorylated AKT; PI3K, phosphoinositide 3-kinase; pGSK3 β , phosphorylated glycogen synthase kinase-3 beta; pTDP-43, phosphorylated TDP43; ROS, reactive oxygen species; SEM, standard error of the mean; TDP-43, transactive response DNA binding protein 43

References: 1. Hultsz D. Am J Manag Care. 2018;24(15):S320-S326. 2. Tortelli R et al. Front Neurol. 2020;11:552295. 3. Cohen T et al. Trends in Molecular Medicine 2011;17(11):659-667. 4. Scotter EL et al. Neurotherapeutics. 2015;12(2):352-363. 5. Berthiaume AA et al. Front Neuro. 2024;18. 6. Ishigaki A et al. J Neuropathol Exper Neurol. 2007;66:1037-1044

Acknowledgments

This study was sponsored and funded by Athira Pharma, Inc. Research support was provided by Neuro-Sys SAS (Gardanne, France) and In Vivex SAS (Lunel, France) funded by Athira Pharma, Inc.


Disclosures

Sherif M Reda, Andrée-Anne Berthiaume, Wei Wu, Kayla N Kleist, Sharay E Setti, Robert W Taylor, Jewel L Johnston, and Kevin J Church are employees and stockholders of Athira Pharma, Inc.

Disclaimer


ATH-1105 is an investigational therapy and has not received FDA approval nor been demonstrated to be safe or effective for any use.

Figure 4. ATH-1105 supports mitochondrial health and reduces caspase-3 activation following glutamate injury




Quantification of (A) functional mitochondria (via MitoTracker) and (B) caspase-3 activation 24h post glutamate challenge expressed as percentage of normal control (100%). Data presented as mean \pm SEM; n = 4-6. Statistical differences were determined by one-way ANOVA followed by Fisher's LSD test versus glutamate control. *p < 0.05, **p < 0.01, ****p < 0.0001

Figure 5. ATH-1105 reduces pTDP-43 and NfL, and improves neuromuscular function in the Prp-TDP43^{A315T} mouse model of ALS



Quantification of (A) sciatic nerve pTDP-43, (B) plasma NfL, and (C) CMAP amplitude after 2 months of treatment. Data presented as mean \pm SEM; n = 10. Statistical significance was determined by 2-way ANOVA with the Dunnett test versus ALS + vehicle. ****p < 0.0001.

Figure 6. Proposed avenues by which ATH-1105 attenuates neurodegenerative mechanisms associated with TDP-43 pathology



Numbers correspond to figure numbers